Factors Associated with the Incidence of COVID-19 Disease in Muang Pattani District, Thailand

Lukman Dunthara, M.Sc.¹, Arinda Ma-A-Lee, Ph.D.¹ Surasak Sangkhathat, M.D., Ph.D.³

¹Department of Mathematics and Computer Science, Faculty of Science and Technology, Prince of Songkla University, Mueang, Pattani 94000, Thailand.

²Department of Biomedical Sciences and Biomedical Engineering, Faculty of Medicine, Prince of Songkla University, Hat Yai, Songkhla 90110, Thailand.

³Translational Medicine Research Center, Faculty of Medicine, Prince of Songkla University, Hat Yai, Songkhla 90110, Thailand. Received 25 August 2023 • Revised 1 September 2023 • Accepted 1 September 2023 • Published online 7 February 2024

Abstract:

Objective: This study aimed to investigate the Coronavirus disease (COVID-19) incidence in Muang Pattani district; from April 2021 and September 2022, Specifically, the main objective was to study the varying distribution by demographic, area and period of COVID-19.

Material and Methods: COVID-19 data, with 15,807 reported cases was obtained from the Muang Pattani District Health Office. A descriptive analysis was used to describe the characteristics of the COVID-19 cases. A negative binomial model was used to investigate COVID-19 incidence.

Results: The overall incidence of COVID-19 was 1,188.9 cases per 100,000 of the population. Females had a higher incidence rate of COVID-19 infection than males. People aged from 20 to 39 years had significantly higher incidence rates than the average, with the highest rate being found in the Barohom sub-district. In 2021, the highest rates were in the month of October, followed by July and August; while March was reported as the highest rate in 2022.

Conclusion: The findings of this study can be used by the public health organizations and others involved in COVID-19 for direct prevention and control in those high-incidence areas. This study has also proven that its findings can be applied to cases related to the prevention programs of other infectious diseases in Thailand.

Keywords: COVID-19 incidence, demographic, negative binomial, Thailand, spatial-temporal

This paper was from the 2nd Annual Health Research International Conference (AHR-iCON, July 19-20, 2023). Contact: Arinda Ma-A-Lee, Ph.D. Department of Mathematics and Computer Science, Faculty of Science and Technology, Prince of Songkla University, Mueang, Pattani 94000, Thailand. E-mail: arinda.m@psu.ac.th

© 2024 JHSMR. Hosted by Prince of Songkla University. All rights reserved. This is an open access article under the CC BY-NC-ND license (http://www.jhsmr.org/index.php/jhsmr/about/editorialPolicies#openAccessPolicy). J Health Sci Med Res doi: 10.31584/jhsmr.20241035 www.jhsmr.org

Introduction

Corona virus disease 19 (COVID-19) is a contagious disease of the respiratory system. It is brought about by an infection of a novel coronavirus strain that was eventually given the name Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) initially identified in Wuhan, China, in December 2019. Later in March 2020, the disease spreaded to over 200 countries, resulting in a total of almost 300 million cases and 5 million deaths¹. The COVID-19 pandemic also caused the SARS-CoV-2 virus to mutate, resulting in the emergence of many new variants. Among Alpha them, Beta, Gamma, Delta, and Omicron were included in the list of variants of concern. The increased rates of transmission, pathogenicity and immune evasion by Omicron set it apart from the initial Wuhan's variant. These variants have triggered a new wave of epidemics worldwide².

The distribution of COVID-19 varies depending on spatial and temporal factors. The population density and distance between each city were found to be highly linked with the number of infected people in a prior study.^{3,4} Furthermore, as new SARS-CoV-2 variants emerged, the characteristics of the disease distribution changed over time. According to the study by Lak et al. (2021)⁵, the demographic composition and physical characteristics (living space and public transport) were key factors explaining the rates of infection and deaths. From previous studies, it was found that demographic factors affecting infection include age groups and gender. Those between the ages of 20 and 49, as well as those 50 and older, were at a comparatively higher risk of infection^{6,7}. Although, the risks for males and females are similar, males had a higher in hospitals mortality rate⁸. Additionally, while COVID-19 vaccination still cannot completely prevent infection, it can reduce the risk of hospitalization and death from infection⁹.

The COVID-19 outbreak in Thailand can be divided into five phases¹⁰. Each wave caused an increase in COVID-19 patients as a result from the emergence of new variants¹⁰. The spread of disease was also due to social activities. In a previous study, the provinces with higher economic status and number of visitors, including Bangkok and Phuket, had a higher number of infected cases¹¹. A study conducted in Thung Yang Daeng district, where the first cluster of COVID-19 was discovered in Pattani province, revealed that contact related to the Islamic missionary (Dawah Tabligh) activities, were at a significantly higher risk of COVID-19 infection than other community members¹². This cluster was the first outbreak in the lower south of Thailand.

COVID-19 studies have been reported in Thailand. However, most of the studies in the southernmost were based on preliminary results. In consequence, the study findings based on the adjusted rate are crucial and more accurate results. Therefore, this research took one crucial step further by applying statistical modeling to the COVID-19 data in order to obtain a more accurate incidence rate for this study. This research aimed to investigate factors associated with the incidence of COVID-19 in the Muang Pattani district.

Material and Methods

Data source and management

The infected patients in this study were Thai people from Pattani that tested positive by RT-PCR. The Muang Pattani District Public Health Office collected all reported cases from April 2021, when the first cases in the district were discovered, until September 2022; the last month before the outbreak was declared endemic. There were 16,997 reported cases. The exclusion criteria were as follows: 1) Patients with other nationalities were not counted, because their population numbers are not reported. Although the population number of migrant workers was variables, it cannot be used due to the number being lower than the reported COVID-19 cases. 2) Patients whose were not fully available study variable. As a result, the total number of infected people stood at 15,807.

Dunthara L, et al.

The dependent variable of this study was the incidence of COVID-19. This study used the population data from the Pattani Provincial Public Health Office, which divides the population by year, gender, age group, and sub-district.

Statistical analysis

In this study, data analysis utilized descriptive statistics analysis tools; such as frequency and percentage, to describe the sample's characteristics. A poisson regression model was later fitted for investigating factors associated with COVID-19 incidence. This model is considered appropriate for modeling count data. However, Poisson models for disease counts are frequently overdispersed; whereby, in such case the negative binomial model is more appropriate. The negative binomial model for incidence rates is an extension of the Poisson model for disease counts that provides for the over-dispersion, which is typical in disease counts.

The plot of residuals against normal quantiles is then used to assess how well a model fits with the data. For the Negative binomial model deviance residuals based on the likelihood are recommended. The plotted residuals should correspond to the normal quantiles by following a straight line on the plot.

The sum contrasts were used to generate confidence intervals for comparing the adjusted incidence rates within each factor in the overall incidence rate. The R program was used for statistical analysis and graphical displays.

Results

Descriptive statistics

Table 1 shows that 16,065 cases of COVID-19 were mostly female (58.5%), within the age groups of 20-29 years (18.49%). Most of them were from the area of Sabarang (20.47%), followed by Bana (19.63%), and Ru Samilae (16.82%), respectively. The highest rates in 2021 were found in the month of October (17.12%), followed by July

(13.31%) and August (13.38%); while in 2022 the highest rates were found in March (17.09%).

Table 1 Characteristics of infected patients (N=16,065 cases)

Variables	Frequency	Percentage
Gender		
Female	9,401	58.52
Male	6,664	41.48
Age group (years)		
0-9	1,861	11.58
10–19	2,120	13.2
20-29	2,971	18.49
30-39	2,931	18.24
40-49	2,309	14.37
50-59	1,949	12.13
60+	1,924	11.98
Sub-district		
Anoru	1,080	6.72
Bana	3,154	19.63
Baraho	756	4.71
Barahom	245	1.53
Chabang Tiko	976	6.08
Kamiyo	355	2.21
Khlong Maning	278	1.73
Paka Harang	563	3.50
Puyut	804	5.00
Ru Samilae	2,702	16.82
Sabarang	3,289	20.47
Talubo	1,050	6.54
Tanyong Lulo	813	5.06
Month-year		
April 2021	53	0.33
May 2021	91	0.57
June 2021	710	4.42
July 2021	2,138	13.31
August 2021	2,134	13.28
September 2021	1,298	8.08
October 2021	2,751	17.12
November 2021	1,813	11.29
December 2021	667	4.15
January 2022	156	0.97
February 2022	937	5.83
March 2022	2,745	17.09
April 2022	329	2.05
May 2022	84	0.52
June2022	30	0.19
July 2022	31	0.19
August 2022	62	0.39
September 2022	36	0.22

Statistical modeling

Figure 1(a) depicts the residual plot for the Poisson model. The model fails due to the appearance of overdispersion, and the chi-square test confirmed that the model is unfit. However, Figure 1(b) shows that the negative binomial model fitted the data well, and the chi-square test indicates the fit is statistically acceptable. The results based on this model are shown in Figure 2.

Figure 2 shows the adjusted COVID-19 incidence rates per 100,000 of the population, with 95% confidence intervals. These confidence intervals used sum contrasts

to compare the COVID-19 incidence rate with the overall mean. This is represented by a red horizontal line, which equals to1188.9 per 100,000. Females had a higher incidence rate than males in addition to the highest incidence rate being found in people aged 20–29, followed by people aged 30–39. Barahom was the sub-district with the significantly highest incidence rate. while the incidence rates below the average were found in Baraho, Chabang Tiko, Klong Maning, and Kamiyo, respectively. The highest incidence rates in 2021 occurred in the month of October, followed by July and August; whereas, in 2022, they occurred in March.



Figure 1 Residual plot for the Poisson model (a) and Negative binomial



Figure 2 95% confidence interval plot of factors associated with COVID-19 incidence rate

Discussion

The findings in this study demonstrated that the four factors; namely: age, district, month, and year, were all associated with the rate of COVID-19 incidence. This study found that females had a higher infection rate than males. This is in contrast to earlier studies, which concluded that there was no difference between the sexes as to the chance of infection¹³. In a study, it was found that females were less susceptible to infection than males, because of the two sexes' different immune responses and between sexes the higher proportion of Angiotensin Converting Enzyme 2 (ACE-2) in males.9 ACE-2 is a receptor protein for the spike protein for the pathogen COVID-19, and infection is brought on by both proteins¹⁴. Hence, this may explain why females have higher infection rates. However, women were more likely than men to visit a doctor¹⁵, which would probably have the same impact on the likelihood to undergo RT-PCR or ATK testing.

This study found that the age groups of 20 and 39 had a higher incidence rate than other groups; especially

in children. This is found to be consistent with the results of the study during the first wave by Jakhmola et al.⁶, which found that those aged 20–49 and those aged 50 and over are at higher risk of contracting COVID–19. This is due to the presence of the ACE-2 gene in the nasal epithelium being the lowest in children under 10 years of age, which then increases with age. However, during the outbreak of the Omicron variants, younger children were more likely to become severely ill than older children16; especially those aged 0–4, most likely due to their ineligibility for vaccination against COVID–19¹⁷.

The map of 13 sub-districts in Muang Pattani district is shown in Figure 3. The map also presents the adjusted COVID-19 incidence rate when compared with the mean incidence; which equals to 1188.9 per 100,000 population. Barahom had a significantly highest incidence rate, while, Baraho, Chabang Tiko, Khlong Maning, and Kamiyo subdistricts had incidence rates around the mean incidence. This might be due to certain commonalities between these sub-districts; for example, their population and housing



Figure 3 A map of sub-district in muang Pattani district with adjusted incidence rate

densities are higher than others. Similarly, to a study by Lak (2021)⁵, the findings revealed that factors contributing to the high number of infected people in urban society in Tehran, Iran, consisted of population density, the density of houses the number of neighborhood shopping and retail centers, and so forth. In addition, a study in China found that the two key factors explaining the spread of the pandemic were human movement and population density¹⁸.

In 2021, the months with the high incidence rates were July-November, which was during the country's fourth wave10. In this study the peak of the incidence was found in October, similarly to neighboring provinces; such as Yala, Narathiwat, and Songkhla. This incidence was found to be contradicting to the overall trend of the country, where a peak of COVID-19 cases was found in August and gradually decreased in October¹⁹. The predominant variant of SARS-CoV 2 in this wave was Delta¹⁰; whereby, it was found that Delta can transmit more efficiently than Alpha and Beta: the dominant variants of the previous waves20. The Delta variant has also been reported to have immune escape capability, reducing the effectiveness of AstraZeneca and Pfizer-BionTech vaccines against infection and hospitalization when compared to the alpha variant²¹⁻²². In 2022, the month with the highest incidence rate in this area was March. This incidence is consistent with the trend of the country; wherein, the peak of COVID-19 cases occurred throughout the outbreak and during the country's fifth wave, January-May 2022, with Omicron as the predominant variant10. A study by Tian et al. (2022)23 found that the Omicron variant had 1.4-3.1 times more infectivity than the Delta variant. This was because of the following reasons: i) the majority were asymptomatic or mild²⁴, and ii) two doses of the AstraZeneca and Pfizer-BionTech vaccines were insufficient to provide adequate protection²⁵ compared to other strains, and more likely to result in re-infection23. The policy of not confirming all possible cases with PCR testing, as well as the widespread use of antigen test kits $(ATK)^{26}$, may have resulted in a lower incidence rate during the fifth wave than in the fourth wave.

Conclusion

This study revealed that demographic, area, and period are all factors associated with COVID-19 incidence. However, suggestions for further studies, in order to get more specific detailed results; especially in other specific areas, should include other potential factors; such as economic status and COVID-19 vaccination. Additionally, advanced infectious models incorporating with spatiotemporal dimensions could be used in further, subsequent research for explanation of the virus mutation and transmission characteristics. The findings of this study can be used by public health organizations and others involved in prevention of COVID-19 and other infectious diseases. Moreover, the statistical methods utilized in this study can be applied to other regions in Thailand and other countries to investigate factors associated with the incidence of COVID-19 or other diseases.

Acknowledgement

We sincerely appreciate the data provided by the Muang Pattani District Public Health Office and the Pattani Provincial Public Health Office. This study was funded by a thesis research grant from the Graduate School, Prince of Songkla University, and tuition fee funding from the Faculty of Science and Technology, Prince of Songkla University, Pattani Campus, Thailand.

Conflict of interest

The authors declare no conflicts of interest.

References

 World Health Organization. Statement on the sixth meeting of the International Health Regulations (2005) Emergency Committee regarding the coronavirus disease (COVID-19) pandemic [homepage on the Internet]. Geneva: WHO; 2021. [cited 2023 Apr 24]. Available from: https://www.who.int/news/item/15-01-2021-statement-on-the-sixth-meeting-of-the-internationalhealth-regulations-(2005)-emergency-committee-regardingthe-coronavirus-disease-(covid-19)-pandemic

- Choi JY, Smith DM. SARS-CoV-2 variants of concern. Yonsei Med J 2021;62:961. doi: org/10.3349/ymj.2021.62.11.961
- Feng Y, Li Q, Tong X, Wang R, Zhai S, Gao C, et al. Spatiotemporal spread pattern of the COVID-19 cases in China. PLoS One 2020;15:e0244351. doi: 10.1371/journal. pone.0244351
- Ruksakulpiwat S, Zhou W, Chiaranai C, Saengchut P, Vonck JE. Age, sex, population density and COVID-19 pandemic in Thailand: A nationwide descriptive correlational study. J Health Sci Med Res 2021;40:281-91. doi: 10.31584/jhsmr.2021836
- Lak A, Sharifi A, Badr S, Zali A, Maher A, Mostafavi E, et al. Spatio-temporal patterns of the COVID-19 pandemic, and place-based influential factors at the neighborhood scale in Tehran. Sustain Cities Soc 2021;72:103034. doi: 10.1016/j. scs.2021.103034
- Jakhmola S, Baral B, Jha HC. A comparative analysis of COVID-19 outbreak on age groups and both the sexes of population from India and other countries. J Infect Dev Ctries 2021;15:333–41. doi: 10.3855/jidc.13698
- Aslam S SM, Suhail K M, Kulkarni A. Clinical and epidemiological characteristics of elderly patients with Coronavirus Disease–19 at a tertiary care center in South India. J Health Sci Med Res 2023;2023963. doi: 10.31584/jhsmr.2023963
- Nguyen NT, Chinn J, De Ferrante M, Kirby KA, Hohmann SF, Amin A. Male gender is a predictor of higher mortality in hospitalized adults with COVID-19. PLoS One 2021;16:e0254066. doi: 10.1371/journal.pone.0254066
- Adamoski D, Baura VA de, Rodrigues AC, Royer CA, Aoki MN, Tschá MK, et al. SARS-CoV-2 Delta and Omicron variants surge in Curitiba, southern Brazil, and its impact on overall COVID-19 lethality. Viruses 2022;14:809. doi: 10.3390/ v14040809
- World Health Organization. WHO Thailand Weekly Situation Update No. 243 [homepage on the Internet]. Geneva: WHO;
 2022. [cited 2023 Apr 24]. Available from: https://cdn.who.int/ media/docs/default-source/searo/thailand/2022_07_27_thasitrep-243-covid-19.pdf?sfvrsn=23bab9e2_1

- Tantrakarnapa K, Bhopdhornangkul B, Nakhaapakorn K. Influencing factors of COVID-19 spreading: a case study of Thailand. Z Gesundh Wiss 2022;30:621–7. doi: 10.1007/s10389-020-01329-5
- Phiriyasart F, Chantutanon S, Salaeh F, Roka A, Thepparat T, Kaesaman S, Sangsawang C, et al. Outbreak investigation of coronavirus disease (COVID-19) among Islamic missionaries in southern thailand. Outbreak, surveillance. OSIR 2020;13:48–54.
- Ballering AV, Oertelt-Prigione S, olde Hartman TC, Rosmalen JGM, Boezen M, Mierau JO, et al. Sex and gender-related differences in COVID-19 diagnoses and SARS-CoV-2 testing practices during the first wave of the pandemic: The dutch lifelines COVID-19 cohort study. J Womens Health (Larchmt) 2021;30:1686–92. doi: 10.1089/jwh.2021.0226
- Hu B, Guo H, Zhou P, Shi Z-L. Characteristics of SARS-CoV-2 and COVID-19. Nat Rev Microbiol 2021;19:141–54. doi: 10.1038/ s41579-020-00459-7
- Thompson AE, Anisimowicz Y, Miedema B, Hogg W, Wodchis WP, Aubrey-Bassler K. The influence of gender and other patient characteristics on health care-seeking behaviour: a QUALICOPC study. BMC Fam Pract 2016;17:38. doi: 10.1186/ s12875-016-0440-0
- Butt AA, Dargham SR, Loka S, Shaik RM, Chemaitelly H, Tang P, et al. Coronavirus disease 2019 disease severity in children infected with the Omicron variant. Clin Infect Dis 2022;75:e361–7. doi: 10.1093/cid/ciac275
- Marks KJ, Whitaker M, Anglin O, Milucky J, Patel K, Pham H, et al. Hospitalizations of children and adolescents with laboratoryconfirmed COVID-19 — COVID-NET, 14 states, July 2021– January 2022. MMWR Morb Mortal Wkly Rep 2022;71:271–8. doi: 10.15585/mmwr.mm7107e4
- Han Y, Yang L, Jia K, Li J, Feng S, Chen W, et al. Spatial distribution characteristics of the COVID-19 pandemic in Beijing and its relationship with environmental factors. Sci Total Environ 2021;761:144257. doi: 10.1016/j.scitotenv.2020.144257
- World Health Organization. WHO Thailand Weekly Situation Update No. 206 [homepage on the Internet]. Geneva: WHO;
 2021. [cited 2023 Jul 10]. Available from: https://cdn.who.int/ media/docs/default-source/searo/thailand/2021_10_21_engsitrep-206-covid19.pdf?sfvrsn=9ad4bfdf_5
- 20. Campbell F, Archer B, Laurenson-Schafer H, Jinnai Y, Konings F, Batra N, et al. Increased transmissibility and global

spread of SARS-CoV-2 variants of concern as at June 2021. Euro Surveill 2021;26:2100509. doi: 10.2807/1560-7917. es.2021.26.24.2100509

- Sheikh A, McMenamin J, Taylor B, Robertson C. SARS-CoV-2 Delta VOC in Scotland: demographics, risk of hospital admission, and vaccine effectiveness. Lancet 2021; doi: 10.1016/s0140-6736(21)01358-1
- 22. Twohig KA, Nyberg T, Zaidi A, Thelwall S, Sinnathamby MA, Aliabadi S, et al. Hospital admission and emergency care attendance risk for SARS-CoV-2 delta (B.1.617.2) compared with alpha (B.1.1.7) variants of concern: a cohort study. Lancet Infect Dis 2022;22:35–42. doi: 10.1016/s1473-3099(21)00475-8
- 23. Tian D, Sun Y, Xu H, Ye Q. The emergence and epidemic characteristics of the highly mutated SARS-CoV-2 Omicron

variant. J Med Virol 2022;94:2376-83. doi: 10.1002/jmv.27643

- Dyer O. Covid–19: South Africa's surge in cases deepens alarm over omicron variant. BMJ 2021;375:n3013. doi: 10.1136/bmj. n3013
- Andrews N, Stowe J, Kirsebom F, Toffa S, Rickeard T, Gallagher E, et al. Covid–19 vaccine effectiveness against the omicron (B.1.1.529) variant. N Engl J Med 2022;386:1532–46. doi: 10.1056/NEJMoa2119451
- World Health Organization. COVID-19 Situation, Thailand [homepage on the Internet]. Geneva: WHO; 2022. [cited 2023 Jul 10]. Available from: https://cdn.who.int/media/docs/defaultsource/searo/thailand/2022_03_30_tha-sitrep-229-covid-19. pdf?sfvrsn=49c43172_5