Original Article



Study Protocol for a Single-center Randomized Clinical Trial: Evaluating the Effectiveness of Combined Therapy versus Conventional Therapy Approaches to Improve Physical & Behavioral Status among Children with Autism Spectrum Disorder

Sherajul Haque, MPT-Paediatrics¹, Abid Hasan Khan, MPT-MSKM (Fellow)¹, Feroz Kabir, Ph.D.¹, Sharmila Jahan, M.Phill¹, Zahid Hossain, Ph.D. (Fellow)^{1,2}, Ehsanur Rahman, M.Sc.PT¹, Nupur Akter, MPT-Gynecology & Women's Health (Fellow)¹, K M Amran Hossain, M.Sc.PT¹

¹Department of Physiotherapy & Rehabilitation, Faculty of Health Science, Jashore University of Science & Technology, Jashore 7408, Bangladesh.

²International Ph.D Program in Gerontology and Long-Term Care, College of Nursing, Taipei Medical University, Taipei 110, Taiwan.

Received 28 December 2024 • Accepted 16 January 2025 • Published online 14 May 2025

Abstract:

Objective: Currently, traditional Autism Spectrum Disorder (ASD) treatments include occupational therapy and speech and language therapy (SLT), however, physiotherapy has little evidence of efficacy. Hence, this study aims to compare physiotherapy, occupational therapy as well as speech and language therapy as need-based interventions for ASD patients with physical and behavioral deficits.

Material and Methods: An assessor and participant blinded randomized clinical trial (RCT) was designed for 70 cases diagnosed with ASD, according to Diagnostic and Statistical Manual of Mental Disorders (DSM) V, from July 2024 to April 2025, with each being randomly assigned to either Group A or Group B. Participants of Group A will receive physiotherapy plus occupational therapy with SLT and group B will receive only occupational therapy plus speech and language therapy for 4 weeks. The outcome of this study will be physical status determined by modified SF-36 scores and behavioral status via GARS-3 scores. The study site will be Proyash (an Institute of Special Education), Jashore Area, Bangladesh. Study analysis will be conducted according to the nature of the data, and will include intention-to- treat analysis.

Contact: Dr. K M Amran Hossain, M.Sc.PT

Department of Physiotherapy & Rehabilitation, Faculty of Health Science, Jashore University of Science & Technology, Jashore 7408, Bangladesh.

E-mail: kma.hossain@just.edu.bd

© 2025 JHSMR. Hosted by Prince of Songkla University. All rights reserved.

This is an open access article under the CC BY-NC-ND license
(http://www.lhsmr.org/index.php/lhsmr/about/editorialPolicies#openAccessPolicy).

J Health Sci Med Res 2026;44(1):e20251207 doi: 10.31584/jhsmr.20251207 www.jhsmr.org **Results:** The results of this study are expected to determine that comprehensive multidisciplinary practices of ASD can advocate the benefits for children and adolescents with ASD, the parents, other stakeholders and special schools throughout Bangladesh.

Conclusion: This study will address the research gap on the impact of the outcome of physiotherapy interventions, along with occupational therapy with SLT, on behavioral and physical status for the participants of ASD.

Keywords: Autism Spectrum Disorder (ASD), behavioral status, physical and emotional health, rehabilitation

Introduction

Autism Spectrum Disorder (ASD) is one of the fastest-evolving cognitive disorders in Bangladesh that affects social interaction and basic behavior¹. Although it can be diagnosed at any age, symptoms of autism usually appear in the initial two years of life². Diagnostic Criteria of Autism in Bangladesh is according to the Directorate General of Health Services guidelines; however, symptoms of autism vary from child to child. Generally, these can be divided into three categories: social barriers, difficulties/ barriers in establishing communication, and repetitive and stereotyped. Both social and communication skills of children with autism do not develop normally. Normally, parents are the first to notice abnormal behavior in a child³. According to the Neuro-Developmental Disabilities Protection Trust law diagnosis criteria, among which the presence of the symptoms above are mentioned, includes the following clauses: (a), (b) and (c) is certain, and clauses (d), (e), (f), (g), (h), (i), (j) and (k), if one or more of the symptoms are observed, they will then be considered as a disabled person having features of ASD; namely:- (a) limitations in verbal or non-verbal communication, (b) limitation of social and interpersonal behavior, communication and imaginative activities, (c) repetition of similar or limited acts or conduct, (d) greater or lesser sensitivity than others in hearing, sight, smell, taste, touch, pain, balance and movement, (e) intellectual disability or any other handicap or disability, (f) extraordinary ability in one or more specific subjects

and inequality of development in the same person, (g) not making or keeping less direct eye contact with others, (h) excessive excitability, excitement or incoherent crying, (i) unusual physical gestures, (j) excessive tendency to follow the same routine, and (k) any other feature prescribed by the Government, from time to time; by Gazette Notification⁴. Children with ASD often exhibit repetitive or restricted behaviors in interaction, language, and communication as well as a variety of other social problems⁵. People with ASD come from diverse socioeconomic backgrounds, races, and nations⁶. Presently, the incidence of ASD in Bangladesh is progressively increasing⁷. Nonetheless, adequate epidemiological data concerning the total prevalence of autism cases in Bangladesh is lacking. According to the Bangabandhu Sheikh Mujib Medical University Center for Child Neurodevelopment and Autism in Bangladesh, 7.5 out of every 10,000 children are diagnosed with autism8. The Institute for Pediatric Neurodisorders and Autism (IPNA) is a tertiary care institution, where the majority of children with ASD in the metropolitan area come for treatment9. A 2013 national study of 7,200 people in Bangladesh found that 1.5 out of every 1,000 children had ASD³.

The rate was highest in Dhaka Metropolis (30 per 10,000 children), while the rate was lower in rural areas (7 per 10,000 children)¹⁰. In 2016, the National Center for Health Statistics estimated that one in 36 children had been diagnosed with Autism. A significant disparity in the ratio of males to girls has also been observed: notably 4:1³. The

severity of ASD is categorized into three levels; wherein, Level 1, indicates moderate autism that necessitates assistance; Level 2, which demands substantial support; and Level 3, representing the most severe type of autism that requires extensive aid11. Nonetheless, a child's growth, development, and contextual variations might affect the intensity of social skills and behaviors¹². Moreover, children with ASD face an elevated risk of comorbidities compared to their typically developing peers¹⁰. Individuals with ASD exhibit lower levels of activity compared to their normally developing counterparts. Research indicates that social and emotional deficiencies, along with related comorbidities, particularly in children with ASD, may be intensified by insufficient physical activity and delayed motor skills and fitness¹³. Physical, behavioral, and emotional challenges associated with ASD may be intensified, significantly adversely affecting an individual's well-being. Presently, there is no recognized therapy for ASD. Considered the gold standard for treating core symptoms, behavioral therapy works best when started early in life¹⁴. However, very few people benefit from this initial, intensive behavioral therapy, as most people with ASD require lifelong supportive care¹⁵. The current conventional therapy approach for autism in Bangladesh consists of only speech and language therapy and occupational therapy, wherein physiotherapy is not a treatment of choice. Research has shown that physical activity helps to improve social interaction, reduce aggressive behavior, and reduce stereotypical behavior in children with ASD16. ASD children enhance social skills and restricted and repetitive behavior patterns through integrated physical training¹⁷. Additionally, exercise improves various comorbidities and symptoms including alleviating physical motor deficits¹⁸.

At present, no study has compared conventional ASD therapies (occupational therapy with speech and language therapy) with combined therapies (physiotherapy,

occupational therapy with speech and language therapy) to improve children's behavioral, physical, and emotional well-being. Hence, combination therapy must be evaluated to improve Bangladeshi autistic children's physical and behavioral parameters. This study will compare integrated treatments, using physiotherapy, occupational therapy, and speech and language therapy, versus need-based treatment, such as occupational therapy and speech and language therapy, for children with ASD.

Material and Methods

Study design

This study will be an assessor and participant-blinded Randomized Clinical Trial. The Standard Protocol Items: Interventional Trials (SPIRIT) – 2013 criteria will be followed to ensure transparency of experimental studies (Table 1).

Sample size calculation

The sample size was determined utilizing ClinCalc software, concentrating on the primary outcome assessed by the Gilliam Autism Rating Scale: Second Edition (GARS – 2)¹⁹. The sample size was established according to the expected minimum clinically significant differences (MCID) of GARS-2: calculated at 4.7±1.7 (on a 0–10 Hiva scale derived from a 0–126 GARS-2 score). The full sample size is 66, as calculated based on a 25% minimum clinical improvement, a 1:1 enrollment ratio, 80% power, and an alpha value of 0.05²⁰. To ensure safety, we will recruit 70 children with ASD, which will then be divided into two groups: 35 in each group.

Recruitment and randomization

The study's population will consist of children with ASD at Proyash, an institute of special education at Jashore Cantonment, from July 2024 to April 2025. Participant

Table 1 Standard procedure of the study protocols, according to SPIRIT guidelines

Time point	Enrolment	Allocation	Post-allocation		
	-T ₁		T,	T ₂	
Enrolment Eligibility screen Informed consent Demographic asso Group allocation Intervention PT OT SLT Assessment GARS-3		x x	X	X X X	
SF-36					

PT=physiotherapy, OT=occupational therapy, SLT=speech and language therapy, GARS-3=Gilliam Autism Rating Scale, T₀=group allocation, T₁=baseline before the intervention, T₂=measurement taken in 3-months after T₂

selection and eligibility screening will be voluntary. At the entrance, patients will be diagnosed with ASD using the Directorate General of Health Services and Neuro-Developmental Disabilities Protection Trust guidelines. They will be randomly assigned to groups after screening. Next, participants will be divided into two groups using sequentially numbered envelopes at 1:1 ratio. The "Rand" function of Microsoft Excel 2021 will perform secret allocation.

Study procedure

After the study registration, researchers will recruit willing participants. Parents will be invited to complete a demographic survey to screen "ASD" cases. Computergenerated random selection will choose 70 autistic children for investigation. An assessor, blinded to the group assignment, will pre-test and screen on the first day. One postgraduate physiotherapists, an occupational therapists, and a speech-language pathologists will treat the patient together. Accessories vary by treatment setup. This study will follow CONSORT guidelines.

Study setting

The study will be conducted at Proyash (Institute of Special Education), Jashore Area, Bangladesh. Study participants will be recruited randomly from the Outpatient Service Unit of Proyash (Institute of Special Education), Jashore Area, Bangladesh.

Eligibility criteria

Participants will be included in this study if they: (1) are between the ages of 3 and 22¹⁹, (2) were diagnosed as children with ASD, and, based on DSM-V criteria²¹, (3) are willing to participate. Participants will not be included if: (1) the respondent consents but drops out within the first week of enrollment; (2) have other neurodevelopmental disorders; such as cerebral palsy, Down syndrome and are children with intellectual disabilities; (3) have any recent (within the past six months) self-reported or parent-reported muscle or joint injury or disease²².

Outcome measurements

Socio-demographic and clinical information

Sociodemographic information, such as age, gender, mother's education, living area, monthly family income, and clinical information, like BMI, birth complication, type of delivery, first child or not, and so forth, will be gathered by a structured questionnaire²³.

Diagnosis of ASD

Screening of children with autism will be performed according to the DSM-5 committee diagnostic criteria²². The three main defining characteristics of ASD are: limited and repetitive activity patterns, deficits in verbal and nonverbal communication, and disruption of social relationships. These basic characteristics can be present irrespective of socio-economic background, caste, nationality, or culture. However, because individuals with ASD are often unique, one trait may be more common than another^{24,25}. Consequently, the current clinical diagnosis of ASD is based on behavioral assessment according to the DSM-5 criteria, published by the American Psychological Association²⁶. The first diagnostic category of the DSM-5 includes deficits in social communication and interpersonal interaction.

According to the third type, symptoms should be visible from childhood; however, they may fully manifest when societal expectations exceed an individual's potential. Symptoms must disrupt regular activity. The diagnosing psychologist offers a severity level of 1 to 3, based on social interactions, restricted interests, and repetitive behavioral patterns. Severity level 3 signifies that this domain is creating significant hindrances for the user²⁷. This instrument has approximately 98.5% sensitivity and 92% specificity in the diagnosis of ASD²².

Behavioral assessment

ASD children's behavior will be assessed using the Gilliam Autism Rating Scale (GARS-3); third edition. The

58-item scale measures Restricted/Repetitive Behaviors (13), Social Interaction (14), Social Communication (9), and Emotional Responses (8). Cognitive Style (seven measures) and Maladaptive Speech (seven items) are used for talkative youngsters. The child's current actions are evaluated with a four-point Likert scale: "not at all like the individual" (0), "not much like the individual" (1), "somewhat like the individual" (2), and "very much like the individual" (3). If an object's evaluation is ambiguous, the youngster should be monitored for an extended period, or prior observational data should be utilized. The average Cronbach's alphas for the Autism Index 4 and 6 are 0.94 and 0.93 for the four and six subscales, respectively¹⁹.

The test-retest reliabilities of 122 people examined twice in two weeks ranged from 0.76 to 0.87 throughout the subscales, with the Autism Index 4 and 6 scoring 0.90. An exploratory factor study verified DSM-5 diagnostic indicators. The scale's predictive validity showed that the autism index scores could distinguish children with ASD from typically developing children, with a sensitivity of 0.96 (4 subscales) and 0.95 (6 subscales) in addition to a specificity of 0.95 and 0.97. However, when comparing ASD children to other developmental illnesses, the index has lower specificity (0.78 for 4 subscales, 0.84 for 6 subscales)²⁰.

Physical functioning and emotional health

A self-administered questionnaire consisting of 36 items, wherein the SF-36 yields eight scores. Four of these relate to physical health: general health, role limitations due to physical problems, physical discomfort, and physical functioning. The remaining characteristics are: vitality, mental health, role limitations due to affective illness, and social functioning. Each scale is rated on a range from 0 to 100. If a person gets a score of 100 on physical functioning, role limitation due to physical or mental problems, physical discomfort, or social functioning, it means no limitation or disability²⁸. A modified SF-36 questionnaire will be employed

to evaluate the physical functioning and emotional well-being of children with ASD, focusing on activity limitations, physical health issues, and mental health concerns²⁹. We will utilize three domains of the SF-36: activity limits, physical health issues, and mental health concerns, via the study's aims.

Intervention

Group A (Physiotherapy and occupational therapy with speech-language therapy)

Physiotherapy interventions:

Group A will receive physiotherapy, such as balance training on land and in water, coordination training, hippotherapy or therapeutic horseback riding, proprioception training, enhance motor function and lower limb strength, training with aerobic exercise to enhance sleep, motor skills, mental state, and muscle strengthening to enhance higher executive performance 30,31. Physiotherapy interventions will be conducted daily for 30 minutes in 1.5-hour daily sessions.

Occupational therapy interventions:

Sensory integration will be customized to each child's unique needs, using the ten basic treatment methods outlined in the Credibility Tool. Some of these methods are: (a) arranging space to improve communication; (b) being physically safe; (c) providing alternatives to sensory input; (d) attaining and preserving suitable levels of stimulation; (e) providing qualitative challenges by changing activities; (f) ensuring the success of activities; (g) providing advice on behavior self-regulation; (h) establishing an enjoyable atmosphere; (i) selecting activities together; and (j) promoting a therapeutic partnership. These ten topics will be divided into three main groups: (1) creating sensory opportunities for the child by changing the environment during treatment; (2) providing common challenges and promoting adaptive responses; and (3) building a rapport between the therapist and the child³². Play therapy, balancing training, skill development drills, ball games, pegboard games, and

sensory integration exercises are some other interventions that incorporate gross motor function through exercise³³. Also, cognitive behavioral therapy and community mobility or travel training have been included³⁴. Occupational therapy interventions will be conducted daily for 30 minutes in 1.5–hour daily sessions.

Speech and language therapy interventions:

Group A will undergo Speech and Language Therapy (SLT); incorporating methods such as: meanings Applied Behavior Analysis (ABA)³⁵, Applied Behavior Consequences (ABC), Relationship Development Intervention (RDI), Cognitive Behavior Intervention (CBI), meanings Picture Exchange Communication System (PECS); Augmentative and Alternative Communication (AAC) as well as Social communication strategies; such as Social Stories, and visual schedules^{36,37}. Each session included 30 minutes of SLT within a 1.5-hour daily therapy schedule.

Group B (Speech-language therapy and occupational therapy)

Group B will receive occupational therapy with speech and language therapy, as per the above protocol, for 1.5 hours per session: each therapy intervention will continue for 45 minutes; however, they will not receive any physiotherapy.

Participants in both groups will be provided 1 hour and 30 minutes of each session, per day for 3 days a week, for up to 4 weeks. After 4 weeks, the assessor will obtain the post-test data.

Minimization of bias and the blinding process

The study will use a double-blind procedure to guarantee that neither participants, nor assessors are aware of the group allocations. Randomization will be performed by an independent party not associated with

the project. Participants would be kept unaware of other group therapies. Distinct responsibilities will be assigned to the examiner, therapy provider, monitoring group, and trial supervisor to enable effective working combinations. The research group will not participate in the study's execution.

Data monitoring

Two people, not directly participating in the trial, will constitute the monitoring team. Their duties shall include: overseeing the intervention procedure, side effects, and participant group registration. Furthermore, they will examine the data and perform an interim analysis. If the Institutional Review Board (IRB) implements any modifications to the research protocols or interventions, the lead investigator shall notify them.

Safety measures and managing adverse effects

While the treatment is predicted to have a small number of side effects, the monitoring group will document any unforeseen incidents occurring during or after the treatment and quickly inform the appropriate specialists. The therapist will capture these occurrences in their notes and communicate them to the lead investigator.

Data audit and management:

Patient codes will be used to fill out a printed, anonymous questionnaire. A volunteer team from the Master of Physiotherapy program at Jashore University of Science & Technology's Department of Physiotherapy & Rehabilitation will enter, audit, and analyze trial data under the supervision of a faculty member not involved in trial operations. The corresponding author of Jashore University of Science & Technology's Department of Physiotherapy and Rehabilitation will keep the paper questionnaire and digital dataset secret.

Data analysis

Statistics will be performed with IBM SPSS 26. Data distribution will be examined using Kolmogorov–Smirnov and Shapiro–Wilk normality tests. For continuous and discrete data, descriptive statistics will provide the mean, median, standard deviation, or interquartile range (IQR); whereas, categorical data will use frequencies and percentages. We will examine baseline compatibility and group differences following the test using the independent t–test or Mann–Whitney U test, depending on data distribution. Within–group comparisons will use Wilcoxon signed–rank or paired sample t–tests for paired variables. Intention–to–treat analysis accounts for one–month treatment dropouts, and statistical significance is defined as a p–value<0.05.

Ethical issues and informed consent

The Institutional Review Board (IRB) of the Physiotherapy and Rehabilitation Department from Jashore University of Science & Technology approved the trial on May 20, 2024 (PTR-JUST/IRB/2024/05/222403). The Clinical Trial Registry India (CTRI) prospectively registered the experiment on July 8, 2024 (CTRI/2024/07/070209). The administrative authorities of Proyash Institute of Special Education, Jashore will provide authorization for the study. The researcher will follow the Helsinki Declaration ethics. Participants will give written informed permission before enrolling and can withdraw at any time without affecting therapy. Only the lead investigator, data auditors, and authors will have access to the anonymised trial dataset. Contractually, investigators that independently examine and document patient data cannot receive or share the final dataset. In case of failure, post-trial therapy will be given.

Results

We anticipate that the complete multidisciplinary practice of ASD may be advocated throughout this

study. Additionally, this will be beneficial to children and adolescents ASD as well as to the parents and special schools in Bangladesh. The results of this study will further define certain stakeholders.

Discussion

This study aims to evaluate the efficacy of combination therapy in enhancing the physical and behavioral status of children with ASD, addressing a worldwide concern among this population. Multiple studies have discussed various therapeutic modalities, including physiotherapy, occupational therapy, and speech and language therapy, to address the physical and behavioral conditions of children with autism. Nevertheless, no research has directly contrasted the effectiveness of physiotherapy with occupational and speech and language treatment approaches. Due to insufficient supporting evidence, a rigorously planned, randomized clinical trial is essential to assist therapeutic doctors in clinical decision—making in this aspect.

This focus on reflecting actual clinical practice increases the study's significance and applicability. Ultimately, the findings will enhance the comprehension of effective interventions for children with ASD, addressing a problem that profoundly affects neurodevelopmental diseases, such as autism spectrum disorder, worldwide. This research addresses a significant gap in existing knowledge by comparing physiotherapy therapies with occupational and speech–language therapy, offering vital guidance to doctors making treatment options for ASD. The study's findings will aid in establishing an effective, comprehensive strategy for addressing the severity of autism, regulating stereotyped behaviors, and reducing potential health risks associated with malnutrition in children with ASD.

Conclusion

This study can aid in the formulation of comprehensive treatment guidelines for ASD in Bangladesh. The study's

strength lies in its representative sampling derived from a population-based cross-sectional home survey, ensuring methodological rigor. Consequently, the study findings are restricted to individuals with ASD living in Jashore City, Bangladesh. Future multicenter trials throughout all eight divisions of Bangladesh may yield results with enhanced external validity.

Acknowledgement

Authors acknowledge the contribution of Col. Monsur Ahmed (BA- 6483, PSC), Chief Co-Ordinator; Mina Sultana, Principal (Acting), and Mst. Sumaya Sultana, Clinical Physiotherapist, Proyash Jashore Area, for their cordial support in the study process.

Funding sources

University Grant Commission (UGC) of Bangladesh granted a partial fund through the university and department. Grant number: 24–FoHS–1.

Conflict of interest

The authors declared that they have no known competing financial interests or personal relationships that could have appeared to influence the work that will be reported in this paper.

References

- Lyall K, Croen L, Daniels J, Fallin MD, Ladd-Acosta C, Lee BK, et al. The changing epidemiology of autism spectrum disorders. Annu Rev Public Health 2017;38:81–102.
- Long D, Yang T, Chen J, Dai Y, Chen L, Jia F, et al. Age of Diagnosis and demographic factors associated with autism spectrum disorders in chinese children: a multi-center survey. Neuropsychiatr Dis Treat 2022;18:3055-65.
- Akhter S, Hussain AE, Shefa J, Kundu GK, Rahman F, Biswas A. Prevalence of autism spectrum disorder (ASD) among the children aged 18–36 months in a rural community of Bangladesh: a cross sectional study. F1000Res 2018;7:424.

- Khatun R, Bin Siddique MK, Khatun MR, Benzir M, Islam MR, Ahmed S, et al. Nutritional status of children with neurodevelopmental disorders: a cross-sectional study at a tertiary-level hospital in northern Bangladesh. BMC Nutr 2024;10:61.
- Tian J, Gao X, Yang L. Repetitive restricted behaviors in autism spectrum disorder: From mechanism to development of therapeutics. Front Neurosci 2022;16:780407.
- Li YA, Chen ZJ, Li XD, Gu MH, Xia N, Gong C, et al. Epidemiology of autism spectrum disorders: Global burden of disease 2019 and bibliometric analysis of risk factors. Front Pediatr 2022;10:972809.
- Williams DL, Siegel M, Mazefsky CA. Problem behaviors in autism spectrum disorder: association with verbal ability and adapting/coping skills. J Autism Dev Disord 2018;48:3668–77.
- Zablotsky B, Black LI, Blumberg SJ. Estimated prevalence of children with diagnosed developmental disabilities in the United States, 2014–2016. NCHS Data Brief 2017;291:1–8.
- Loomes R, Hull L, Mandy WP. What is the male-to-female ratio in autism spectrum disorder? A systematic review and metaanalysis. J Am Acad Child Adolesc Psychiatry 2017;56:466-74.
- Must A, Curtin C, Hubbard K, Sikich L, Bedford J, Bandini L.
 Obesity prevention for children with developmental disabilities.
 Curr Obes Rep 2014;3:156–70.
- Burns CO, Matson JL. An evaluation of the clinical application of the DSM-5 for the diagnosis of autism spectrum disorder. Expert Rev Neurother 2017;17:909-17.
- Broder-Fingert S, Brazauskas K, Lindgren K, Iannuzzi D, Van Cleave J. Prevalence of overweight and obesity in a large clinical sample of children with autism. Acad Pediatr 2014;14:408-14.
- Gharehgazlou A, Vandewouw M, Ziolkowski J, Wong J, Crosbie J, Schachar R, et al. Cortical gyrification morphology in ASD and ADHD: implication for further similarities or disorder–specific features? Cereb Cortex 2022;32:2332–42.
- Jackson SL, Abel EA, Reimer S, McPartland JC. Brief report: a specialized fitness program for individuals with autism spectrum disorder benefits physical, behavioral, and emotional outcomes.
 J Autism Dev Disord 2024;54:2402–10.
- 15. Batey CA, Missiuna CA, Timmons BW, Hay JA, Faught BE, Cairney J. Self-efficacy toward physical activity and the physical activity behavior of children with and without Developmental Coordination Disorder. Hum Mov Sci 2014;36:258-71.

- 16. Alhowikan AM, Elamin NE, Aldayel SS, AlSiddiqi SA, Alrowais FS, Hassan WM, et al. Children with autism spectrum disorder exhibit elevated physical activity and reduced sedentary behavior. Brain Sci 2023;13:1575.
- Oriel KN, George CL, Peckus R, Semon A. The effects of aerobic exercise on academic engagement in young children with autism spectrum disorder. Pediatr Phys Ther 2011;23:187– 93
- Ferreira JP, Ghiarone T, Cabral Junior CR, Furtado GE, Moreira Carvalho H, Machado-Rodrigues AM, et al. Effects of physical exercise on the stereotyped behavior of children with autism spectrum disorders. Medicina 2019;55:685.
- 19. Samadi SA, Biçak CA, Noori H, Abdalla B, Abdullah A, Ahmed L. Autism spectrum disorder diagnostic criteria changes and impacts on the diagnostic scales-utility of the 2nd and 3rd versions of the Gilliam Autism Rating Scale (GARS). Brain Sci 2022;12:537.
- 20. Akter R, Urme NA, Hossain KA, Hossain T, Ahammad S, Yeasmin MH, et al. Protocol for a randomized clinical trial comparing the efficacy of Structured Diet (SD) and Regular Therapy (RT) for adolescents with malnutrition having Autism Spectrum Disorder (ASD). PLoS One 2023;18:e0292326.
- Huerta M, Bishop SL, Duncan A, Hus V, Lord C. Application of DSM-5 criteria for autism spectrum disorder to three samples of children with DSM-IV diagnoses of pervasive developmental disorders. Am J Psychiatry 2012;169:1056-64.
- 22. Wiggins LD, Rice CE, Barger B, Soke GN, Lee LC, Moody E, et al. DSM-5 criteria for autism spectrum disorder maximizes diagnostic sensitivity and specificity in preschool children. Soc Psychiatr Epidemiol 2019;54:693-701.
- 23. Faruk MO, Rahman MS, Rana MS, Mahmud S, Al-Neyma M, Karim MS, et al. Socioeconomic and demographic risk factors of autism spectrum disorder among children and adolescents in Bangladesh: evidence from a cross-sectional study in 2022. PLoS One 2023;18:e0289220.
- 24. Moulton E, Bradbury K, Barton M, Fein D. Factor analysis of the childhood autism rating scale in a sample of two year olds with an autism spectrum disorder. J Autism Dev Disord 2019;49:2733-46.
- 25. Lord C, Elsabbagh M, Baird G, Veenstra-Vanderweele J. Autism spectrum disorder. Lancet 2018;392:508-20.
- 26. Khan NZ, Gallo LA, Arghir A, Budisteanu B, Budisteanu M,

- Dobrescu I, et al. Autism and the grand challenges in global mental health. Autism Res 2012;5:156-9.
- Gulati S, Kaushik JS, Saini L, Sondhi V, Madaan P, Arora NK, et al. Development and validation of DSM-5 based diagnostic tool for children with Autism Spectrum Disorder. PLoS One 2019:14:e0213242.
- Lins L, Carvalho FM. SF-36 total score as a single measure of health-related quality of life: Scoping review. SAGE Open Med 2016;4:2050312116671725.
- 29. Fong DY, Chan BK, Li S, Wan CH, Kazis LE. Average and individual differences between the 12-item MOS Short-form Health Survey version 2 (SF-12 V. 2) and the veterans RAND 12-item Health Survey (VR-12) in the Chinese population. Health Qual Life Outcomes 2022;20:102.
- 30. Amonkar N, Su WC, Bhat AN, Srinivasan SM. Effects of creative movement therapies on social communication, behavioralaffective, sensorimotor, cognitive, and functional participation skills of individuals with autism spectrum disorder: a systematic review. Front Psychiatry 2021;12:722874.
- Peters BC, Pan Z, Christensen H, Gabriels RL. Self-regulation mediates therapeutic horseback riding social functioning outcomes in youth with autism spectrum disorder. Front Pediatr 2022;10:884054.

- 32. Belaiba M, Laatar R, Borji R, Ben Salem A, Sahli S, Rebai H. Time Limited Benefits of Physical and Proprioceptive Training on Physical Fitness Components in Children With Autism Spectrum Disorders. Percept Mot Skills 2024;131:785–804.
- 33. Suárez-Manzano S, Ruiz-Ariza A, de Loureiro NE, Martínez-López EJ. Effects of physical activity on cognition, behavior, and motor skills in youth with autism spectrum disorder: a systematic review of intervention studies. Behav Sci 2024;14:330.
- 34. Smith E, Constantin A, Johnson H, Brosnan M. Digitally-mediated social stories support children on the autism spectrum adapting to a change in a 'real-world'context. J Autism Dev Disord 2021;51:514-26.
- Kuhaneck H, Spitzer SL, Bodison SC. A systematic review of interventions to improve the occupation of play in children with autism. OTJR (Thorofare N J) 2020;40:83–98.
- 36. Storch EA, Schneider SC, De Nadai AS, Selles RR, McBride NM, Grebe SC, et al. A pilot study of family-based exposure-focused treatment for youth with autism spectrum disorder and anxiety. Child Psychiatry Hum Dev 2020;51:209–19.
- Kersten ML, Coxon K, Lee H, Wilson NJ. Developing community mobility and driving with youth on the autism spectrum: a psychosocial perspective. Scand J Occup Ther 2021;28:91–6.