

Factors Influencing Retention of Medical Graduates in the Health System: A Study from The First Collaborative Project to Increase Production of Rural Doctors (CPIRD) in Thailand

Watcharin Chirdchim, M.D., Ph.D.¹, Sirapong Weawseetong, M.D.²,
Nutthaeak Thangwatthanarat, M.D.², Koramit Navacharoenwong, M.D.²

¹Department of Obstetrics and Gynecology, Prapokklao Hospital, Mueang, Chanthaburi 22000, Thailand.

²Medical Education Center Prapokklao Hospital, Mueang, Chanthaburi 22000, Thailand.

Received 14 November 2024 • Revised 9 January 2025 • Accepted 16 January 2025 • Published online 22 May 2025

Abstract:

Objective: To identify the factors affecting doctor retention in the public health system.

Material and Methods: This cross-sectional study surveyed 426 doctors who graduated from Prapokklao Hospital (1976–2018). Data collected included: (1) Basic information (gender, age, resignation age, service length, medical program); (2) Personal factors (income, work hours, shifts, patient load, education, position); (3) Psychological factors (job satisfaction, pride); (4) External factors (relationships, welfare, culture, risks). Statistical analysis involved t-tests, Chi-square tests, and logistic regression to adjust for biases.

Results: Continuing in a residency program significantly increased retention in the public health service (odds ratio=7.14, 95% confidence interval=4.72, 10.81, p-value<0.05).

Conclusion: Residency training is a key factor for enhancing doctor retention in the public health system.

Keywords: CPIRD, doctor retention, resignation

Contact: Watcharin Chirdchim, M.D., Ph.D.
Department of Obstetrics and Gynecology, Prapokklao Hospital, Mueang,
Chanthaburi 22000, Thailand.
E-mail: watcharin.ch@cpird.in.th

J Health Sci Med Res
doi: 10.31584/jhsmr.20251213
www.jhsmr.org

© 2025 JHSMR. Hosted by Prince of Songkla University. All rights reserved.
This is an open access article under the CC BY-NC-ND license
(<http://www.jhsmr.org/index.php/jhsmr/about/editorialPolicies#openAccessPolicy>).

Introduction

From 1960 to 1975, there was a significant and persistent shortage of doctors in the public health system in Thailand. During this period, a large number of Thai doctors resigned from government positions and chose to work abroad. As a result, only approximately 300 doctors remained to serve in rural areas^{1,2}. In 1976, negotiations between the Ministry of Public Health and Chulalongkorn University led to the establishment of a project aimed at producing doctors for rural areas. This project focused on educating and addressing problems in rural settings. Prapokklao Hospital was the first clinical medical education center in Thailand to produce doctors under the medical education for students in rural area project (MESRAP), a collaboration between Chulalongkorn University and the Ministry of Public Health. Initially, the project was called “The MESRAP,” launched in 1976. Later, in 1994, the project was renamed and expanded into 2 programs: the collaborative project to increase production of rural doctors (CPIRD) and the one district one doctor (ODOD) program³.

Despite efforts through the CPIRD to reduce resignations and increase retention in the public health system, the challenge of keeping doctors remains⁴⁻⁶. Even with increased doctor production, retention issues persist⁷. Over 40 years, Prapokklao Hospital has trained more than 800 doctors for rural areas, emphasizing real-world problem-solving in these settings. However, no study has specifically examined retention and resignation rates from these programs. This research aimed to identify the factors influencing doctor retention and resignation, guiding program improvements for sustainable development in line with World Health Organization standards¹³.

Material and Methods

This cross-sectional descriptive study analyzed factors influencing doctor retention and resignation in the

MESRAP, ODOD, and CPIRD programs at Prapokklao Hospital. Ethical approval was granted (Approval Number: CTIRECT 04/65) covering the period from September 8, 2022, to September 7, 2023. Participants were informed about the study and provided written consent. Data were collected from November 1, 2022, to August 31, 2023, with 337 questionnaires completed. The study population included 853 doctors who graduated from Prapokklao Hospital's rural doctor program between 1976 and 2018.

The sample size was calculated using Taro Yamane's formula (1973) at a 95% confidence level, requiring 273 participants. Systematic sampling selected 426 doctors. Inclusion criteria were doctors who graduated from the CPIRD, ODOD, and MESRAP programs at Prapokklao Hospital (1976–2018) and consented to participate. Exclusions included non-graduates, those who declined participation, deceased individuals, unreachable contacts, or retirees. Data were collected through phone interviews, covering: 1) Basic information (age, gender, service duration, program type), 2) Personal factors (income, workload, education, position), 3) Psychological factors (job satisfaction), and 4) External factors (work environment, relationships, risks).

Data analysis was conducted using STATA software. Descriptive statistics included mean and standard deviation for quantitative data, and percentages for qualitative data. Inferential analysis used independent t-tests for quantitative data and chi-square tests for qualitative data. Binary logistic regression identified factor relationships, while survival analysis and Kaplan–Meier comparisons evaluated retention rates, with significance set at $p\text{-value} < 0.05$ and a 95% confidence interval (95% CI).

Results

The study included 853 doctors who graduated from the rural doctor program (1976–2018). From this group, 426

were randomly selected for surveys, with 67 excluded based on the criteria, resulting in a final sample of 337 doctors. Ages ranged from 27 to 62 years, averaging 41.8 years. After excluding retirees, 262 doctors (81.1%) remained in the public health system, while 61 (18.9%) had left, as illustrated in Figure 1.

The basic information reveals key findings indicating significant differences in age between doctors who stayed in the system (40.2 ± 7.3 years) and those who resigned (44.6 ± 9.63 years). The average resignation age was 31.8 ± 6.9 years. Length of service and gender significantly influenced retention, with an equal male-to-female ratio (50%) among those who stayed. In the MESRAP program, 70.53% remained, while 29.47% resigned. In the ODO program, 85.71% stayed and 14.29% left. For the CPIRD program, 86.8% remained and 13.2% resigned, as shown in Table 1.

Section 2 analysis shows significant differences: doctors in the system earn less but work more hours daily and weekly (40.3 vs. 36.0) compared to those who resigned, with fewer shifts (2.5 vs. 3.2). More doctors who stayed pursued training (46.6% residency, 22.9% fellowships) than those who resigned (21.3% and 8.2%). Most who resigned had no additional training (70.5% vs. 30.5%). Resigned doctors were mainly practicing (52.5%) and senior physicians (23.0%), while those who stayed were specialists (21.0%), experts (17.2%), and highly qualified physicians (11.5%). The differences are statistically significant ($p\text{-value} < 0.05$), as shown in Table 2. Job satisfaction between the 2 groups shows no significant differences. Most reported being moderately happy (68.32% retained, 67.21% resigned) and moderately proud (64.50% retained, 68.85% resigned). Satisfaction with supervisors (83.61% resigned vs. 73.66% retained) and welfare (73.77% resigned vs. 60.31% retained) was slightly higher among those who resigned. Satisfaction

with corporate culture (86.89% resigned vs. 72.14% retained) and risk (82.25% resigned vs. 77.48% retained) also followed this trend. Despite these variations, the differences aren't statistically significant enough to impact retention or resignation decisions, as shown in Table 3.

Table 4 identifies the key factors influencing doctors' decisions to stay or leave the public health system, analyzed through univariable and multivariable methods: age, years of service, gender, income, daily hours, training, and position. Doctors who stay tend to be younger, have fewer years of service, a lower income, work longer hours, and hold higher positions. Continuing education, especially residency, increases the likelihood of staying, while female doctors are more likely to resign. Satisfaction with supervisors, colleagues, welfare, and culture had no significant impact. Notably, doctors who stay earn less ($p\text{-value} < 0.05$) but are up to 7 times more likely to pursue further education ($p\text{-value} < 0.05$). Researchers used survival analysis and Kaplan-Meier with log-rank tests to further analyze the factors impacting retention. Results showed that continuing education significantly influences the decision to resign from the public health system ($p\text{-value} = 0.000$, $p\text{-value} \leq 0.05$), as shown in Table 5.

Using the Kaplan-Meier method, we found that the highest likelihood of resigning occurs within the first 10 years, especially in the first 5 years of service. Retention improves after 10 years, but the likelihood of resigning rises again after 20 years of service, as shown in Figure 2.

In the first 10 years, retention rates dropped most for doctors without residency training, followed by those with residency, and least for those with advanced residency. By the 30th year, retention was around 60% for non-residency doctors, 70% for residency-trained, and 75% for those with advanced residency, as shown in Figure 3.

Table 1 Personal information of doctors between those who remain in the system and those who have resigned

Parameter	Retain (n=262) mean (\pm S.D.)	Not-retain (n=61) mean (\pm S.D.)	p-value
Age (years)	40.2 (\pm 7.3)	44.6 (\pm 9.6)	<0.05
Age in work	40.2 (\pm 7.3)	31.8 (\pm 6.9)	<0.05
Year work	16.1 (\pm 7.5)	7.5 (\pm 7.2)	<0.05
Gender (n, %)			<0.05
Male	131.0 (50.0)	20.0 (32.79)	
Female	131.0 (50.0)	41.0 (67.22)	
Programs (n, %)			<0.05
ODOD	12.0 (4.58)	2.0 (3.28)	
CPIRD	171.0 (65.27)	26.0 (42.62)	
MESRAP	79.0 (30.15)	33.0 (54.10)	

S.D.=standard deviation, ODOD=one district one doctor, CPIRD=collaborative project to increase production of rural doctors, MESRAP=medical education for students in rural area project

Table 2 Compensation, work intensity, continuing education and current position of doctors between those who remain in the system and those who have resigned

Work intensity	Retain (n=262) mean (\pm S.D.)	Not-retain (n=61) mean (\pm S.D.)	p-value
Income/month (Baht)	70,385.4 (\pm 22,626.83)	78,437.5 (\pm 24,498.29)	0.0026
Daywork (hours/day)	7.4 (\pm 1.6)	6.8 (\pm 2.6)	0.037
Wage/week (shifts)	2.5 (\pm 1.8)	3.2 (\pm 3.8)	0.015
Patients/day	44.3 (\pm 21.3)	44.0 (\pm 22.7)	0.7315
Working hour/week	40.3 (\pm 10.6)	36.0 (\pm 17.5)	0.0220
Continuing education			<0.05
No training	29 (11.07)	18 (29.51)	
Resident	129 (49.24)	27 (44.26)	
Fellow	104 (39.69)	16 (26.23)	
Current position			<0.05
Practicing physician	20 (8.16)	20 (40.00)	
Senior physician	74 (30.20)	14 (28.00)	
Specialist	124 (50.61)	16 (32.00)	
Expert	25 (10.20)	0 (0.00)	
Highly qualified physician	2 (0.82)	0 (0.00)	

S.D.=standard deviation

Table 3 Job satisfaction of doctors between those who remain in the system and those who have resigned

Parameter	Retain (n=262, %)	Not-retain (n=61, %)	p-value
Happiness			0.295
Not happy	0 (0.00)	1 (1.64)	
Slightly happy	5 (1.91)	1 (1.64)	
Moderately happy	179 (68.32)	41 (67.21)	
Very happy	71 (27.10)	18 (29.51)	
Extremely happy	7 (2.67)	0 (0.00)	
Dignity			0.711
Not proud	1 (0.38)	0 (0.00)	
Slightly proud	8 (3.05)	0 (0.00)	
Moderately proud	169 (64.50)	42 (68.85)	
Very proud	75 (28.63)	18 (29.51)	
Extremely proud	9 (3.44)	1 (1.64)	
Supervisor			0.089
Not satisfied	0 (0.00)	1 (1.64)	
Slightly satisfied	13 (4.96)	3 (4.92)	
Moderately satisfied	193 (73.66)	51 (83.61)	
Very satisfied	49 (18.70)	5 (8.20)	
Extremely satisfied	7 (2.67)	1 (1.64)	
Risk			0.351
Not satisfied	1 (0.38)	1 (1.64)	
Slightly satisfied	6 (2.29)	1 (1.64)	
Moderately satisfied	203 (77.48)	52 (82.25)	
Very satisfied	50 (19.08)	7 (11.48)	
Extremely satisfied	2 (0.76)	0 (0.00)	
Colleague			0.127
Not satisfied	0 (0.00)	1 (1.64)	
Slightly satisfied	5 (1.91)	3 (4.92)	
Moderately satisfied	182 (69.47)	43 (70.49)	
Very satisfied	69 (26.34)	12 (19.67)	
Extremely satisfied	6 (2.29)	2 (3.28)	
Welfare			0.188
Not satisfied	1 (0.38)	1 (1.64)	
Slightly satisfied	8 (3.05)	1 (1.64)	
Moderately satisfied	158 (60.31)	45 (73.77)	
Very satisfied	84 (32.06)	13 (21.31)	
Extremely satisfied	11 (4.20)	1 (1.64)	
Corporate Culture			0.056
Not satisfied	1 (0.38)	1 (1.64)	
Slightly satisfied	6 (2.29)	1 (1.64)	
Moderately satisfied	189 (72.14)	53 (86.89)	
Very satisfied	60 (22.90)	6 (9.84)	
Extremely satisfied	6 (2.29)	0 (0.00)	

Table 4 Univariable and multivariable odds ratios for factors influencing retention and resignation of doctors

Parameter	Retain (n=262)	Not-retain (n=61)	p-value	Univariable odds ratio	p-value	Multivariable odds ratio	p-value
Age	40.2 (±7.3)	44.6 (±9.6)	<0.05	0.93	<0.05	-	-
Age in work	40.2 (±7.36)	31.8 (±6.86)	<0.05	1.22	<0.05	1.07 (0.97, 1.18)	0.394
Years of service	16.1 (±7.5)	7.5 (±7.2)	<0.05	1.22	<0.05	1.13 (0.97, 1.32)	0.143
Gender (n, %)			<0.05	4.88	<0.05	0.57 (0.35, 0.93)	0.213
Male	131 (50.0)	20 (32.79)					
Female	131 (50.0)	41 (67.22)	<0.05			-	-
Project (n, %)							
ODOD	12 (4.58)	2 (3.28)		0			
CPIRD	171 (65.27)	26 (42.62)		1.09	0.908		
MESRAP	79 (30.15)	33 (54.10)		0.40	0.246		
Income/month (Baht)	70,385.4 (±22,626.8)	78,437.5 (±24,498.2)	<0.05	1.00	<0.05	0.99 (0.97, 1.01)	<0.05
Daywork (hours/day)	7.4 (±1.6)	6.8 (±2.6)	<0.05	1.16	<0.05	1.37 (1.02, 1.84)	0.200
Wage/week (shifts)	2.5 (±1.8)	3.2 (±3.8)	<0.05	0.90	0.111	-	-
Working hours/week	40.3 (±10.6)	36.0 (±17.5)	<0.05	1.03	<0.05	1.05 (0.83, 1.33)	0.724
Training (n, %)			<0.05				
No training	29 (11.07)	18 (29.51)		0		0	
Resident	129 (49.24)	27 (44.26)	<0.05	2.94	<0.05	7.14 (5.32, 9.58)	<0.05
Fellow	104 (39.69)	16 (26.23)	<0.05	4	<0.05	3.33 (2.25, 4.93)	0.108
Position (n, %)			<0.05				
Practicing physician	20 (8.16)	20 (40.00)		0		0	
Senior physician	74 (30.20)	14 (28.00)	<0.05	5.26	<0.05	2.63 (1.78, 3.89)	0.165
Specialist	124 (50.61)	16 (32.00)	<0.05	7.69	<0.05	3.22 (2.09, 4.96)	0.145
Expert	25 (10.20)	0 (0.00)	-	-	-	-	-
Highly qualified	2 (0.82)	0 (0.00)	-	-	-	-	-

ODOD=one district one doctor, CPIRD=collaborative project to increase production of rural doctors, MESRAP=medical education for students in rural area project




Figure 1 study flow





Figure 2 Comparison of survival functions in the public health system with the variable of years of service (Years)



Green line (Top): Doctors who did not pursue residency training, red line (Middle): Doctors who completed residency training, blue line (Bottom): Doctors who pursued advanced residency training, Analysis time (years)

Figure 3 Comparison of survival functions in the public health system with the variable of continuing education (Years)

Discussion

This study surveyed 426 doctors, achieving a participation rate of 37% (323 participants out of 853), which exceeded the required sample size. Among these participants, 262 remain in the public health system, while 61 have resigned. All participants completed the questionnaire, ensuring comprehensive data collection. Initial analyses identified several significant factors associated with retention in the public health system, which were further explored using logistic regression analysis.

One key finding was the variation in resignation rates across different programs. The MESRAP program exhibited the highest resignation rate, followed by ODO, with CPIRD demonstrating the lowest resignation rate. These findings align with previous studies showing that the CPIRD and ODO programs achieve higher retention rates compared to regular programs⁴. Younger doctors were more likely to remain in the public health system, with older doctors more frequently resigning. This trend corresponds with the transition from the MESRAP to CPIRD and ODO programs, which have been bolstered by increased compensation for rural doctors.

Another factor was the age at the final year of service. On average, doctors who resigned were 31 years old in their final year of service, coinciding with the completion of residency and service obligations. This is consistent with earlier studies suggesting that some doctors resign to pursue further education or specialized training⁴⁻⁷. Gender also played a role, with female doctors exhibiting higher resignation rates than their male counterparts. This may be attributed to the requirements of the CPIRD and ODO programs, which often necessitate service in remote hospitals⁸.

Income levels emerged as another significant factor. Doctors who resigned reported higher average incomes than those who remained. However, this difference is likely due to transitions to private hospitals rather than public hospitals

offering competitive salaries⁸. Prior research highlights that factors like pride, compensation, welfare, and family support significantly influence retention, particularly in rural areas⁹. Working hours also appeared to influence retention. Doctors who remained in the public health system reported working more daily and weekly hours than those who resigned. Interestingly, those who resigned took more weekly shifts, possibly to compensate for lower workloads per shift in private settings compared to their public duties⁸.

Opportunities for continuing education strongly influenced retention. Logistic regression analysis revealed that doctors with access to further education were 7 times more likely to remain in the public health system⁹⁻¹⁰. This aligns with studies identifying educational opportunities as critical for extending tenure in the public health system and highlights the role of career advancement in influencing doctors' decisions. Additionally, past research has identified that one of the key reasons for resigning from the public health system is to pursue further education¹¹.

The current position of doctors was another determinant of retention. Doctors who resigned were mostly practicing physicians, whereas those who remained included a higher proportion of senior physicians and specialists. Career advancement opportunities, therefore, appear to play a pivotal role in retention, as corroborated by previous studies¹⁰.

Contrary to expectations, job satisfaction and pride in work did not show significant differences between the 2 groups. These findings diverge from earlier research that emphasized the importance of happiness and pride in retaining healthcare professionals^{1,9,10,12}. Similarly, no significant differences were observed in factors such as relationships with colleagues and supervisors or organizational culture. Despite this, prior research suggests that organizational policies and commitment can influence retention⁹.

Finally, family emerged as a decisive factor in retention decisions, with its influence surpassing other

organizational factors⁸. This underscores the importance of personal and familial considerations in shaping the career choices of doctors within the public health system.

When analyzing retention rates in the public health system using survival analysis and the log-rank test, considering opportunities for further education, the results align with the earlier discussion.

Limitations and recommendations

Retrospective nature: this descriptive study relies on historical data, with some information dating back over 10 years. Although participants provided complete information, some were uncertain about the accuracy of their responses due to the time elapsed since the events.

Survey design: during the design of the survey, questions were tested on doctors who had graduated over 10 years ago. Some personal questions were difficult for respondents to recall and answer. As a result, certain potentially beneficial questions had to be omitted. These included monthly income at the time of resignation, job position at the time of resignation, workload at the time of resignation, daily and weekly working hours at the time of resignation, spouse's domicile, spouse's income, and trends in further education decisions.

Lack of subgroup analysis: researchers did not perform subgroup analyses, such as categorizing responses by different time periods, for ease of calculation. This omission could result in confounding factors related to the trends and influences of different eras, which could impact retention and resignation decisions.

Sample size and depth of analysis: the large sample size prevented in-depth individual interviews. Consequently, the analysis might not fully explore all aspects and reasons behind each factor. The researchers had to rely on explanations and results from previous studies to interpret the findings.

Conclusion

The opportunity for further education significantly impacts retention, indicating that career advancement, continuing education, promotion, salary increases during education, or other incentives during education may increase retention in the public health system.

References

1. Virasombat N. Physician retention in community hospitals of Thailand: a case study of changwat Nakhon Ratchasima. *J Health Sci Thai* 2018;18:526–36.
2. Virasombat N. Physician retention in community hospitals of Thailand [dissertation]. Bangkok: National Institute of Development Administration; 2009.
3. The collaborative project to increase production of rural doctor. History of CPIRD [homepage on the Internet]. Nonthaburi: The collaborative project to increase production of rural doctor 2019 [cited 2021 Dec 10]. Available from: <https://www.cpird.in.th/index.php/index/site/1>
4. Pagaiya N, Kongkam L, Worarat W, Sriratana S, Wongwinyou K. Rural retention of medical graduates trained by the collaborative project to increase rural doctors (CPIRD). *J Health Syst Res* 2012;6:219–27.
5. Pagaiya N, Sriratana S, Wongwinyou K, Lapkom C, Worarat W. Impacts of financial measures on retention of doctors in rural and public health facilities. *J Health Syst Res* 2012;6:228–35.
6. Pagaiya N, Tangcharoensathien V, Kasemsup V, Chantanir A, Sriratana S, Wongwinyou K. Attitude and rural job choices of newly graduated doctors. *J Health Syst Res* 2012;6:30–9.
7. Churproong S, Uengarporn N. Affecting factors for retention in health system of new doctors from Suranareeuniversity of technology. Bangkok: National Research Council of Thailand; 2016.
8. Sikheawsukwongkot IP. Factors influencing turnover of physicians in Thai public hospitals. *J Lang Relig Cult* 2016;5:183–209.
9. Surakijboworn J. Factors affected physician maintaining in community hospital: case study of state doctors in Pathumthani community hospital [dissertation]. Bangkok: Thammasat University; 2014.

10. Martnok T. Maintenance and turnover of radiological physician in hospitals under department of medical services [dissertation]. Bangkok: Thammasat University; 2017.
11. Arora R, Chamnan P, Nitiapinyasakul A, Lertsukprasert S. Retention of doctors in rural health services in Thailand: impact of a national collaborative approach. *Rural Remote Health* 2017;17:4344.
12. Pruksananonda K, Phonphatkul W, Jalayongjea K, Rattanakhuha L, Dulyakasem K, Phannitamai W. Factors affecting physicians' turnover from public hospitals in Thailand. *Thai J Public Adm* 2004;2:48–87.
13. Department of Economic and Social Affairs. Gold standard for the global goals – a best practice standard to quantify, certify and maximize impact to the SDGs. New York: United Nations; 2018.